The Genetic and Molecular Basis of Speech & Stuttering

Maddy Sanz



For much of the twentieth century, stuttering was widely misunderstood. Explanations often focused on psychological causes such as anxiety, nervousness, or personality traits. Modern research has significantly reshaped this view. Today, stuttering is increasingly understood as a neurodevelopmental condition influenced in part by genetic factors that affect how speech-related brain systems develop and function (Kraft & Yairi, 2012; Neef & Chang, 2024).

Evidence for a genetic contribution first came from family studies. Researchers observed that stuttering often occurs in multiple members of the same family, suggesting that inherited factors may influence susceptibility. Large epidemiological studies estimate that 50–70% of individuals who stutter report a family history of the condition (Yairi & Ambrose, 2013; Kraft & Yairi, 2012). Twin studies provide further support for this conclusion. Identical twins, who share nearly all of their genetic material, show significantly higher concordance rates for stuttering than fraternal twins, indicating that genetic inheritance contributes meaningfully to the disorder (Fagnani et al., 2011).

Although these studies established that genetics plays a role, identifying specific genes involved in stuttering has been a challenge. That changed in 2010, however, when researchers discovered mutations in several genes associated with persistent developmental stuttering. In a study published in the *New England Journal of Medicine*, Kang et al. (2010) identified mutations in three genes, GNPTAB, GNPTG, and NAGPA, in individuals with persistent stuttering. These genes that were implicated were surprising because they are not directly involved in language or motor control. They participate instead in a fundamental cellular process. They participate in the intracellular trafficking of enzymes to lysosomes.



GNPTAB, GNPTG, and NAGPA genes. These genes are implicated in the neurobiology of stuttering & primarily involved in the intracellular trafficking of enzymes to lysosomes.

Lysosomes are cellular structures responsible for breaking down and recycling molecules inside the cell. In order for lysosomes to function properly, specialized enzymes must be transported to them through a targeting system that labels these enzymes with a molecular signal known as mannose-6-phosphate. The proteins encoded by the genes GNPTAB, GNPTG, and NAGPA help generate and process this signal, ensuring that lysosomal enzymes reach the correct cellular destination (Kang et al., 2010; Kang & Drayna, 2012). In simple terms, these genes help the cell determine where certain enzymes should be sent. When the targeting process is disrupted, lysosomal enzymes may not reach their proper location, which can alter cellular maintenance processes and protein turnover. Severe disruptions in this pathway are known to cause rare metabolic disorders such as mucolipidosis. However, the mutations associated with stuttering appear to produce much milder functional changes, rather than complete loss of enzyme activity (Kang & Drayna, 2012; Frigerio-Domingues & Drayna, 2017). This raises an important question: if these genes are involved in basic cellular maintenance, why would their mutation appear primarily as a disorder of speech fluency?



The lysosome. Lysosomes degrade and recycle damaged organelles, misfolded proteins, and excess hormones.

One important factor of these genes' role in stuttering is mutation severity. In the severe lysosomal storage disorders associated with GNPTAB and GNPTG, both copies of the gene are typically damaged by mutations that drastically reduce enzyme function, such as mutations that produce truncated proteins. In contrast, the variants identified in individuals who stutter are often missense mutations, which alter the protein but do not completely eliminate its function. As a result, lysosomal trafficking may be only partially impaired rather than globally disrupted (Kang & Drayna, 2012).

Another factor is cell-type vulnerability. Even though lysosomal trafficking occurs in all cells, mild disruptions in this system may affect certain cell types more than others. Researchers have proposed that subtle trafficking deficits could disproportionately affect a small population of cells that play a critical role in the neural circuits responsible for speech (Frigerio-Domingues & Drayna, 2017). Recent animal research has provided important insight into this possibility. In a study published in the *Proceedings of the National Academy of Sciences*, Han et al. (2019) engineered human

stuttering-associated mutations in the GNPTAB gene into mice. These mice displayed abnormalities in vocalization patterns that resembled key aspects of human stuttering. Surprisingly, the most prominent cellular abnormalities were not found in neurons themselves but in astrocytes, a type of glial cell that supports neural function.

Astrocytes play essential roles in maintaining the brain's cellular environment. They regulate neurotransmitter levels, help maintain ion balance, support synapse development, and contribute to the health of white matter pathways. Han and colleagues found that astrocytes in the corpus callosum, the major fiber tract connecting the brain's hemispheres, showed structural abnormalities in mice GNPTAB mutated mice. When the gene was selectively disrupted in astrocytes, similar vocalization abnormalities emerged, suggesting that astrocyte dysfunction may play a role in the neural mechanisms underlying stuttering (Han et al., 2019). This finding is important because it provides a possible link between lysosomal trafficking defects and the development of speech-related brain circuits. If astrocytes or other support cells function less efficiently during brain development, this could subtly influence how neural networks responsible for speech timing and coordination mature.



An astrocyte, one of the most important neuroglial cells needed for normal neuron function. Astrocytes manage the microenvironment of neurons that is needed for optimal activity.

Additional research suggests that the genes associated with stuttering may also intersect with broader metabolic processes in the brain. For example, a neuroimaging genetics study by Chow et al. (2020) found that expression patterns of two stuttering-related genes, GNPTG and NAGPA, were spatially correlated with brain regions showing structural differences in children with persistent stuttering. The genes associated with these brain regions were enriched for pathways involved in energy metabolism, including glycolysis and oxidative metabolism. These findings raise the possibility that lysosomal trafficking genes may influence brain development through their effects on cellular metabolism and energy regulation.

Further evidence for the importance of intracellular trafficking comes from the discovery of another gene associated with stuttering, a gene called AP4E1. This gene encodes a component of the AP-4 adaptor complex, which is involved in sorting proteins within the cell's Golgi apparatus. Although AP4E1 does not participate directly in the mannose-6-phosphate targeting pathway, it is involved in the broader system of intracellular protein trafficking. Rare variants in this gene have been found at higher frequencies in individuals with persistent stuttering, reinforcing the idea that disruptions in cellular trafficking pathways may contribute to the disorder (Raza et al., 2015).

Taken together, these discoveries suggest that stuttering may arise from subtle disruptions in cellular systems that support neural development rather than from abnormalities in a single "speech gene." Mild impairments in intracellular trafficking may affect how certain brain cells function during development, particularly those involved in maintaining the neural circuits responsible for speech timing, motor control, and auditory feedback.

The genetics of stuttering, however, are complex and remain an active area of research. The currently known genes explain only a small proportion of persistent stuttering cases, and many additional genetic factors remain undiscovered. Nevertheless, the discovery of lysosomal trafficking genes associated with stuttering has transformed scientific understanding of the disorder. These findings demonstrate that stuttering has clear biological foundations and highlight the importance of studying cellular

mechanisms, neural development, and brain networks together. As research continues, genetic discoveries will provide new insights into how speech networks develop and why disruptions in these systems can affect the fluency of spoken language and manifest itself into speech disorders.

References

Fagnani, C., Fibiger, S., Skytthe, A., & Hjelmborg, J. (2011). Heritability and environmental effects for self-reported periods with stuttering: A twin study from Denmark. *Logopedics Phoniatrics Vocology*, 36(3), 114–120.

Frigerio-Domingues, C., & Drayna, D. (2017). Genetic contributions to stuttering: The current evidence. *Molecular Genetics & Genomic Medicine*, 5(2), 95–102.

Han, T.-U., et al. (2019). Human GNPTAB stuttering mutations engineered into mice cause vocalization deficits and astrocyte pathology in the corpus callosum. *Proceedings of the National Academy of Sciences*, 116(35), 17515–17524.

Kang, C., et al. (2010). Mutations in the lysosomal enzyme-targeting pathway and persistent stuttering. *New England Journal of Medicine*, 362(8), 677–685.

Kang, C., & Drayna, D. (2012). A role for inherited metabolic deficits in persistent developmental stuttering. *Molecular Genetics and Metabolism*, 107(3), 276–280.

Kraft, S. J., & Yairi, E. (2012). Genetic bases of stuttering: The state of the art. *Folia Phoniatrica et Logopaedica*, 64(1), 34–47.

Neef, N. E., & Chang, S.-E. (2024). Knowns and unknowns about the neurobiology of stuttering. *PLOS Biology*, 22(2).

Raza, M. H., et al. (2015). Association between rare variants in AP4E1 and persistent stuttering. *American Journal of Human Genetics*, 97(5), 715–725.

Yairi, E., & Ambrose, N. (2013). Epidemiology of stuttering: 21st century advances. *Journal of Fluency Disorders*.