What Happens in the Brain When We Speak? An Overview of the Brain Structures Involved in Speech Production

Maddy Sanz



Speaking often feels effortless. Most people can translate thoughts into words almost instantly. A myriad of networks in the brain interact with each other so that we can answer questions, tell stories, and introduce ourselves without consciously thinking about how speech is produced. Yet from a neuroscience perspective, speech is one of the most complex neural mechanisms that the human brain performs.

Producing even a short phrase requires precise coordination of the language system, motor planning networks, sensory feedback, and timing mechanisms distributed across multiple regions of the brain. Understanding how the brain produces speech serves as a foundation for understanding speech differences like stuttering. Over the past several decades, advances in neuroimaging and neuroscience have allowed researchers to study the brain networks involved in speech production with increasing precision. Studies reveal that speech is not controlled by a single "speech center", but instead emerges from a network of interconnected brain regions working together in fractions of a second.

The process begins when the brain formulates a linguistic message when we decide what we want to say. Regions within the left hemisphere of the brain play an especially crucial role in this development. One of the most widely studied brain regions connected to speech is Broca's area, located in the left inferior frontal gyrus. Discovered in the 19th century as a major modulator in speech, Broca's area is now understood to contribute to the planning and sequencing of speech movements. Modern-day neuroimaging research suggests that this region assists in translating linguistic information into motor programs that guide articulation (Hagoort, 2014). Essentially, before a word can be spoken, the brain first generates a detailed plan required for the movements required to produce that word.



Once a speech plan has been formulated, the brain converts this plan into coordinated muscle movements that physically produce the speech sounds. This process relies on the primary motor cortex, which sends signals via neurotransmitters to muscles of the vocal tract including the tongue, lips, jaw, and larynx. The primary motor cortex, however, is not the only brain region needed to generate the physical movements of muscles for the production of speech. Speech requires extremely precise motor control. Even a single syllable involves multiple muscles activating in carefully timed patterns within fractions of a second. As evidenced by Bouchard et. al (2013), neural activity during speech production has demonstrated that specific regions of the motor cortex correspond to movements of different articulatory structures. Bouchard and colleagues mapped neural activity in the sensorimotor cortex during speech and found that distinct patterns of neural activation correspond to specific articulatory gestures. These findings highlight the extraordinary level of coordination required to produce fluent speech.



Although the motor cortex controls the muscles of speech, other brain structures help regulate the timing and sequencing of these movements, which is arguably just as important as generating the motor signals themselves. The basal ganglia, a group of interconnected nuclei located deep within the brain, have been implicated in this role of

initiating movements and regulating their timing. Speech places particularly high demands on this system as sounds must transition rapidly and smoothly from one to the next. Alm (2004) suggests that abnormalities in the basal ganglia pathways involved in motor timing could contribute to the speech disruptions observed in developmental stuttering.



Another structure involved in coordinating speech movements is the cerebellum which is located at the back of the brain. The cerebellum contributes to motor precision and learning via fine-tuning movements and correcting small errors in motor execution. During speech production, the cerebellum helps ensure that articulatory movements occur with the correct timing and coordination. Research by Ackermann (2008) suggests that cerebellar circuits play an important role in maintaining the smooth execution of speech movements and adapting speech motor patterns over time through motor learning.



Speech production also relies on continuous sensory monitoring. As we speak, the brain listens to our own voice and compares the sounds we produce with the sounds we intended to produce. This process involves the auditory cortex, which provides feedback that allows the brain to detect and correct errors during speech. The interaction between speech motor commands and auditory feedback has been described in the DIVA model of speech production, developed by Tourville and Guenther (2011). According to this model, fluent speech depends on continuous communication between motor planning systems and sensory feedback systems, allowing the brain to make rapid adjustments to speech movements as they occur.

Taken together, these findings illustrate that speech emerges from a complex network of interacting brain systems. Language planning regions in the frontal cortex generate speech plans, motor cortex regions execute articulatory movements, basal ganglia circuits regulate timing and initiation, the cerebellum refines coordination, and auditory systems monitor the resulting speech sounds. Researcher Frank Guenther (2016) describes speech production as one of the most sophisticated sensorimotor behaviors performed by the human brain, requiring continuous coordination between multiple neural systems.

References

Ackermann, H. (2008). Cerebellar contributions to speech production and speech perception. *Trends in Neurosciences*.

Alm, P. (2004). Stuttering and the basal ganglia circuits: A critical review. *Journal of Communication Disorders*.

Bouchard, K. E., et al. (2013). Functional organization of human sensorimotor cortex for speech articulation. *Nature*.

Guenther, F. H. (2016). *Neural Control of Speech*. MIT Press.

Hagoort, P. (2014). Nodes and networks in the neural architecture for language. *Current Opinion in Neurobiology*.

Tourville, J. A., & Guenther, F. H. (2011). The DIVA model: A neural theory of speech acquisition and production. *Language and Cognitive Processes*.